<http://www.diva-portal.org>

Postprint

This is the accepted version of a paper published in *Water Science & Technology*. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the original published paper (version of record):

Nehrenheim, E., Muter, O., Odlaire, M., Rodriguez, A., Cepurnieks, G. et al. (2013)
Toxicity assessment and biodegradation potential of water-soluble sludge containing 2,4,6-trinitrotoluen.
Water Science & Technology, 68(8): 1707-1714
<http://dx.doi.org/10.2166/wst.2013.416>

Access to the published version may require subscription.

N.B. When citing this work, cite the original published paper.

Permanent link to this version:

<http://urn.kb.se/resolve?urn=urn:nbn:se:mdh:diva-22751>

Toxicity assessment and biodegradation potential of water soluble sludge containing 2,4,6-trinitrotoluene

E.Nehrenheim*, O.Muter**, M.Odlare*, A.Rodriguez***, G.Cepurnieks**** and V.Bartkevics****

*School of Sustainable Development of Society and Technology, Mälardalen University, P.O. Box 882, SE-721 23 Västerås, SWEDEN (E-mail: emma.nehrenheim@mdh.se, monica.odlare@mdh.se, adrian.rodriguez@mdh.se)

**Institute of Microbiology & Biotechnology, University of Latvia, 4 Kronvalda blvd., LV-1010 Riga, LATVIA (E-mail: olga.muter@inbox.lv)

***Catalan Institute for Water Research (ICRA), Scientific and Technological Park of the University of Girona, Girona 17003, Spain

****National Diagnostic Centre, 3 Lejupes Str., Riga LV-1076, LATVIA, (E-mail: guntis.cepurnieks@ndc.gov.lv, vadims.bartkevics@ndc.gov.lv)

Abstract The water soluble phase of TNT-containing sludge (SLP) was characterized with regard to TNT concentration, ecotoxicity, and a model biodegradation experiment as evaluation criteria for further development of appropriate treatment technologies. SLP contained 67.8 mg TNT/l. The results of germination and root-elongation tests indicated that SLP had a species-specific phytotoxic effect. The results of a 21 day degradation experiment demonstrated TNT conversion to 4-amino-2,6-DNT and 2-amino-4,6-DNT, with a simultaneous reduction in the total concentration of nitroaromatics. Indigenous microbial activity in the sludge solid phase noticeably increased the total TNT concentration, and was slightly lower than in samples containing only the water soluble phase. Measurement of microbial enzyme activity was used to assess changes in the microbial community during the biodegradation process.

Keywords Biodegradation; enzymatic activity; indigenous microorganisms; phytotoxicity; solubility; TNT

Abbreviations CLE – cabbage leaf extract; DHA – dehydrogenase; FDA – fluorescein diacetate; SLP – sludge liquid phase; SSP – sludge solid phase; TNT – 2,4,6-Trinitrotoluene; 4ADNT – 4-Amino-2,6-Dinitrotiluene; 2ADNT – 2-Amino-4,6-Dinitrotiluene; NACs – nitroaromatic compounds

INTRODUCTION

Explosive compounds enter the environment as a result of demilitarization activities, production of ammunition and explosives, open detonation and burning of explosives at army depots, evaluation facilities, artillery ranges and ordnance disposal sites (Nehrenheim and Odlare, 2010). One of the most widely used explosives is 2,4,6-Trinitrotoluene (TNT). Nitro aromatic compounds (NACs) of this type do not occur naturally and most microorganisms are therefore not adapted to degrade and mineralize these substances. The symmetry around the toluene molecule generates a very stable structure that is extremely resistant to microbial degradation (Ribé et al., 2010). There is an urgent need for a cost effective method for removal of explosive compounds in water and sludge.

Technological approaches for decontamination of TNT- and other NACs-containing wastewaters can be divided into physico-chemical and biological methods, or combinations of these. Physico-chemical treatment of red water via sorption by pine bark (Nehrenheim et al., 2011; Chusova et al., 2012), activated polystyrene microspheres (Meng et al., 2012), activated coke from lignite (Zhang et al., 2011), vacuum distillation (Zhao et al., 2010), oxygen excess at 550°C (Chang et al., 2013), and catalytic degradation using iron nanocatalysts and iron oxide

bentonite nanocomposite (Badawi et al., 2012) are among the methods that have been reported for efficient decontamination of NACs-containing wastewaters.

The incredible versatility inherent in microbial metabolism has meant that explosives have become part of the biogeochemical cycle (Singh et al., 2012). Knowledge of the microbial dynamics that drive TNT biodegradation is limited, particularly in native aquifer sediments where TNT poses a threat to water resources (Fahrenfeld et al., 2012; Harrison and Vane, 2010; Hoffsommer and Rosen, 1972).

The use of slurry and packed-bed bioreactors (Wang et al., 2010; Shen et al., 2009; Mulla et al., 2012), and constructed wetlands (Best et al., 1997; Best et al., 1999; Sikora et al., 1998) have been shown to be effective for NACs-containing wastewater treatment. A mixed microbial population in digested sewage culture degraded 110 mg TNT/l under strict anaerobic conditions in 6 days (Kwon, 2000).

Extracellular enzymes, e.g. from white rot fungi, have been shown to be effective degraders of TNT. However, high production cost inhibits the widespread use of extracellular enzymes for remediation (Rugabber and Talley, 2006).

Ammonium nitrate and chloride as nitrogen sources and molasses as a carbon source were found to be efficient amendments for TNT biodegradation (Yasin et al., 2008). Molasses have been reported as an efficient carbon source for the co-metabolism of explosives in many studies (Clark and Boopathy, 2007; Rodgers and Bunce, 2001; Gerth et al., 2003; Lamichhane et al., 2012). Crude plant extracts (e.g., spinach and parrotfeather) and cabbage leaf extract are effective amendments for the NACs biodegradation process (Medina et al., 2004; Muter et al., 2008, Muter et al., 2012).

Biodegradation of explosives waste is influenced by temperature, oxygen supply, nutrient supply, pH, the availability of the contaminant to the microorganism, the concentration of the waste, and the presence of substances toxic to the microorganisms (e.g. mercury) (US Army Environmental Center, 2000; Wang et al., 2004). Conditions must therefore be carefully controlled in order to gain optimal results in treatment of explosives waste.

The aim of the present study was to study microbiological and ecotoxicological development during TNT-degradation using three different mediums. The specific aim was to characterize the water soluble fraction (SLP) of an demilitarization factory waste sludge, by assessing TNT concentration, toxicity, and biodegradation potential before and after treatment.

METHODS

Characterization of SLP physico-chemical properties

TNT-containing sludge was kindly provided by the former munition utilization plant Nammo Vingåkersverken (Sweden). The concentrations of TNT and its metabolites in the sludge solid phase (SSP) and its water soluble phase (SLP) were measured using a HPLC-DAD (diode array detector) system in gradient mode using the Phenomenex Synergi Hydro 250x4.6 mm column (4 µm particles). Calibration was performed using reference standards (Dr. Ehrenstorfer, Germany).

Ecotoxicological study

Biotests on seed germination and root elongation were performed in triplicate according to EPA 712-C-96-154 guidelines (EPA, 1996) on seeds of cress (*Lepidium sativum*), wheat (*Triticum sp.*), rye (*Secale cereale*), barley (*Hodeum vulgare*), oat (*Avena sativa*), clover (*Trifolium sp.*), alfalfa (*Medicago sativa*), soya bean (*Glycine max*), and radish (*Raphanus sativus*). An SLP dilution factor of 0.5 (100%; 50%; 25%; 12.5%; 0%) was used in the tests.

Microbiological study

The total bacteria counts in SSP and SLP were determined by colony counts from triplicate samples after incubation on Tryptone Glucose Yeast Extract Agar (Sifin, Germany) at +28 °C for 72h.

Biochemical study

Enzyme activities of microorganisms in samples were tested as follows. Dehydrogenase (DHA) activity was determined by reduction of 2-*p*-iodo-3-nitrophenyl-5-phenyltetrazolium chloride (INT) to iodonitrophenylformazan. 50µl of a mixture containing 40mg INT, 1ml 1% glucose, and 20ml 0.25M TRIS was added to 50 µl sample and incubated at +28 °C for 48h. 300 µl extraction solution (ethanol and dimethylformamide, 1:1) was then added to the mixture, vortexed and centrifuged at 5000 rpm after 30 min incubation at ambient temperature. Optical density was measured at 485 nm. Urease activity was determined by the colorimetric method according to NH₃-N formation in urea-amended 0.2 M phosphate buffer pH7.1 (after 48 h incubation at +28 °C) (Kandeler and Gerber, 1988). Concentration of NH₄⁺-N was measured with Nessler reagent by optical density at 425 nm. OD₄₂₅=1 corresponded to a N-NH₄⁺ concentration of 6.0 mg/l. Fluorescein diacetate (FDA) hydrolysis activity was determined by hydrolysis of FDA in 0.06 M phosphate buffer pH7.6 for 60 min at +37 °C (Chen et al., 1988). Optical density was measured at 490 nm. OD₄₉₀=1 corresponded to a FDA concentration of 2.98 mg/l.

Batch experiment

Three variants with different combinations of sludge and inoculum were prepared in duplicate 250 ml Duran bottles (Table 1). The bottles were mixed on a shaking table for 21 days at 100 rpm at ambient temperature. The liquid phase was made up with sterile distilled water and M8* stock solution containing 60 g/l Na₂HPO₄, 30 g/l KH₂PO₄, and 5 g/l NaCl. CLE and molasses were used as amendments. CLE and inoculum (cultivated from the sludge liquid phase) were prepared as described by Muter et al. (2008). Sugar beet molasses contained 42 % of reducing sugars. After the enrichment procedure (48h, 28 °C), inoculum was made up with a cell concentration of 1.0 x 10⁴ CFU/ml using bacteria isolated from sludge on TGA medium. The composition of the enrichment medium for the set L (excluding sludge) is shown in Table 1.

Table 1. The experimental setup with different substrate/inoculum combinations

Set	Sludge liquid phase, ml	Sludge solid phase, g moist w	Water, ml	M8* stock, ml	Molasses, 30 %(w/v), ml	CLE, ml	Inoculum, ml
L*	124	0	38	20	8	10	0
LI**	124	0	18	20	8	10	20
LIS***	124	5	13	20	8	10	20

*L – sludge liquid (water) phase; ** LI - sludge liquid (water) phase + inoculum; *** LIS - sludge liquid (water) phase + inoculum + sludge solid phase.

RESULTS AND DISCUSSION

Physico-chemical characterization of the sludge liquid phase

TNT concentration in air-dried sludge was around 30%. The water soluble phase of the tested sludge contained 67.8 mg TNT/l and no other NACs, i.e. TNT degradation products were found. Current literature values of TNT solubility in water vary widely between 100 and 200 mg/L at room temperature. Solubility in water plays an important role in toxicity and biodegradability of chemicals. Under natural conditions, nitroaromatic compounds have only limited bioavailability due to their low solubility in water. Sludge composition may influence TNT solubility and the presence of heavy metals reduces TNT solubility (Nehrenheim et al 2012). The concentration of explosives in groundwater is affected by the size and surface area of individual explosive particles, the degree of weathering, and the rate of dissolution (Lynch et al., 2002, Speitel et al., 2002). In addition, if two sparingly soluble compounds are present, neither can dissolve to the full extent of its reported aqueous solubility (Clausen et al., 2006). This means that the low TNT concentration in water phase could be due to adsorption on suspended solids or low dissolution rate. Temperature and pH also significantly affect TNT solubility (Lynch et al., 2001; Phelan and Barnett, 2001; McCutcheon and Schnoor, 2003). The pH of the SLP varied between 6.3 - 6.9 during the 21 day incubation which should be considered modest in the effect on TNT solubility but can significantly affect conversion of TNT (Qiao et al., 2010). The redox potential was in the range 55.0 - 65.0 mV.

Ecotoxicological assessment of the sludge liquid phase

Effluents from TNT production have been reported to be toxic to different test organisms, e.g. daphnia, bacteria (Ribeiro et al., 2012). TNT leakage to water, its fate and toxic effect in the environment are important research areas, particularly in the context of marine ammunition dumping sites (Ek et al., 2007). One of the most important characteristics of any toxic compound is its bioavailability. Physico-chemical factors of compounds in contaminated water result in toxic effects on biota.

A comparative study of SLP toxicity was performed with a battery of monocot and dicot plants using seed germination and primary root elongation tests as the main criteria to evaluate toxic effects of TNT under the studied conditions. Inhibition of seed germination by SLP was found to be species-specific. 50% SLP (33.9 mg/l) inhibited germination of cress salad by 50%, but did not influence germination of rape and wheat seeds. However, undiluted SLP inhibited germination of all the tested plants in a species-specific manner (results not shown). Among the tested plants, cress, rye, barley, and radish showed the highest sensitivity to the SLP in the root elongation test (results not shown). Oat, clover and alfalfa were more resistant to the presence of SLP. SLP actually stimulated primary root development in soya bean. SLP exposure thus results in different responses in different plant species.

Large numbers of culturable bacteria were detected in both SSP and SLP, i.e. 2.3×10^6 cfu/g and 5.3×10^3 cfu/ml respectively.

TNT degradation study

A biodegradation study was performed as indicated in Table 1. In this experiment, SLP was amended with mineral salt composition M8*, CLE and molasses. These amendments were previously shown to be efficient for TNT biodegradation (Boopathy et al., 1998, Gerth et al., 2003; Limane et al., 2009). Addition of molasses provided approximately 0.5% reducing sugars (mostly sucrose). Non-diluted cabbage leaf extract also contained reducing sugars (mostly glucose and fructose). Comparative study showed that the concentration of reducing sugars in the CLE prepared from different cultivars was usually in the range 10 - 30 g/100ml, while the total nitrogen and carbon was in the range 0.2 - 1.0 and 0.5 - 1.3 (vol.%) respectively (Grube et al., 2008). This study compared sets with different combinations of sludge liquid phase (L), solid phase (which is water soluble to a limited extent) (LIS) and inoculum originating from the same sludge (LI). As shown in Fig.1, two TNT biodegradation products were formed during incubation, 4-Am-2,6-DNT and 2-Am-4,6-DNT. These compounds are known to be the typical products of microbial biodegradation of TNT (McCormick et al., 1976; Hoffsommer et al., 1978; Schackmann and Muller, 1991; Alvarez et al., 1995; Esteve-Nunez et al., 2001). Addition of inoculum stimulated the TNT degradation process. Thus, after 21 days incubation the total concentration of NACs in media with inoculum was 3.7 mg/l, compared to 12.4 mg/l in media without inoculum (Fig.1). The presence of SSP resulted in an increase in water soluble NACs in the medium within 4 days of incubation. TNT degradation via conversion to other NACs enabled additional solubilisation of TNT. Thus the total concentration of TNT degradation products after 21 days incubation in the presence of SSP was considerably higher compared to the sets without SSP (Fig.1). This may indicate more intensive microbial activity in the presence of SSP. These findings will be examined in greater detail in follow-up studies.

Fig. 1. Degradation of TNT to two recognized degradation metabolites in the sludge liquid phase. Description of the sets is shown in Table 1.

L – sludge liquid (water) phase, LI - sludge liquid (water) phase + inoculum; LIS - sludge liquid (water) phase + inoculum + sludge solid phase.

Microbial activity in the sludge liquid phase with different amendments

DHA activity typically occurs in all intact viable microbial cells. DHA activity therefore correlates to the presence of viable microorganisms and their oxidative capability. In this study, a comparison of DHA activity dynamics showed a wide variation, although activity had a tendency to increase with incubation time in all three sets tested (Fig.2A). After 21 days incubation the DHA activity was highest in set L, i.e. without inoculum. FDA hydrolysis activity was also higher in set L compared to sets LI and LIS (Fig.2B). Addition of SSP resulted in total inhibition of FDA hydrolysis activity on the 21st day of the experiment (Fig.2B). Measurement of FDA hydrolysis has been suggested as an appropriate method in integrated bioecosystem studies because the ubiquitous lipase, protease, and esterase enzymes are involved in the hydrolysis of FDA (Green et al., 2006). Determination of urease activity is based on the formation of ammonium ions in the presence of urea as a substrate. The highest NH_4^+ concentrations were found at the beginning of the experiment in all tested sets. NH_4^+ concentration was lower on the 8th day, and subsequently increased again (Fig.2C). In this study, NH_4^+ concentration was influenced by other factors in addition to urease activity. Biotransformation of TNT may increase NH_4^+ concentration via microbial conversion (Martin et al., 1997, Esteve-Nunez et al., 2001). However, controls lacking urease did not show detectable concentrations of NH_4^+ in the medium.

The number of culturable microorganisms increased in the set with inoculum (LI) from 3.3×10^3 to 2.4×10^5 cfu/ml after 21 days incubation. In turn, addition of SSP to incubation media resulted in a slight decrease in the total count of culturable microorganisms (Fig.2D). As shown in Fig.1, addition of inoculum to liquid medium stimulated TNT degradation. Hence, a decrease of TNT concentration in the SLP can be mostly explained by microbial activity. At the same time, dehydrogenase and FDA hydrolysis activity in the set with inoculum was lower than in the set without inoculum after 21 days incubation. These physico-chemical changes in the medium are most likely due to enhanced i) TNT biotransformation and biodegradation; ii) consumption of nutrients; and iii) accumulation of metabolites, leading to a decrease in microbial enzyme activity in the set with inoculum.

Fig. 2. A, B and C: Changes of the microbial enzyme activity analysed as DHA, FDA hydrolysis, and urease activity during 21 days incubation with SLP. D: Number of colony forming units in the SLP during biodegradation.

L – sludge liquid (water) phase,

LI - sludge liquid (water) phase + inoculum;

LIS - sludge liquid (water) phase + inoculum + sludge solid phase. Error bars represent the standard deviation at 5% level of significance.

CONCLUSIONS

TNT-containing sludge and its water soluble fraction were characterized using different approaches. The main conclusions are:

- The toxicity study revealed a species-specific toxic effect of SLP on higher plants. Cress salad was found to be one of the more sensitive plants in this study.
- Use of indigenous bacteria as inoculum stimulated TNT degradation. Further identification and characterization of bacteria may be an important strategic tool in the context of TNT biodegradation in water.

- Counts of colony forming units and of dehydrogenase and FDA hydrolysis activity dynamics indicated an increase in microbial activity with time in SLP. However, the presence of SSP in liquid medium resulted in strong inhibition of microbial activity on the 21st day of incubation.
- Composition of the liquid medium used in this study was found to be appropriate for TNT-degrading activity of indigenous microorganisms. Further comparative study is needed to optimize the TNT-containing wastewater process with emphasis on its cost efficiency.

Acknowledgements

This work was supported by contract AĪVA 2008/220 from the Ministry of Defence (Republic of Latvia). We also wish to acknowledge our BIOREX co-production partners: Nammo Vingåkersverken AB, Saab Bofors Testcenter AB, Casium AB, Eriksson Patent AB KCEM AB and the Swedish Knowledge Foundation (KKS).

REFERENCES

- Alvarez M. A., Kitts C. L., Botsford J. L. and Unkefer P. J. (1995), *Pseudomonas aeruginosa* strain MA01 aerobically metabolizes the aminodinitrotoluenes produced by 2,4,6-trinitrotoluene nitro group reduction. *Canadian Journal of Microbiology*, **41**, 984–991.
- Badawi, A.M., Ahmed, S.M., Shaban, S.A., Morsy, S.M.I. (2012), Nanotechnology: The next revolution for wastewater treatment (TNT contaminate). *Desalination and Water Treatment*, 40(1-6), 1-6.
- Best EP, Sprecher SL, Larson SL, Fredrickson HL, Bader DF (1999), Environmental behavior of explosives in groundwater from the Milan Army Ammunition Plant in aquatic and wetland plant treatments. Removal, mass balances and fate in groundwater of TNT and RDX. *Chemosphere* 38(14): 3383–3396.
- Best EP, Zappi ME, Fredrickson HL, Sprecher SL, Larson SL, Ochman M (1997), Screening of aquatic and wetland plant species for phytoremediation of explosives-contaminated groundwater from the Iowa Army Ammunition Plant. *Ann. NY Acad. Sci.* 829: 179–194.
- Boopathy R., Manning J. and Kulpa C.F. (1998), A laboratory study of the bioremediation of 2,4,6- TNT contaminated soil using a soil slurry reactor. *Water Environment Research*, **70**(1), 80-86.
- Chang, S.-J. , Liu, Y.-C. , Yang, X.-Q. (2013), Study on explosive wastewater treatment by supercritical water oxidation (Conference Paper). 2nd International Conference on Energy, Environment and Sustainable Development, EESD. *Advanced Materials Research* 610-613, 1934-1938.
- Chen W., Hoitink A.J., Schmitthenner A.F. and Tuovinen O.H. (1988), The role of microbial activity in suspension of damping-off caused by *Pythium ultimum*. *Phytopathology*, **78**, 314-322.
- Chusova O., Nehrenheim E., Odlare M (2012) Dynamic adsorption of TNT-contaminated industrial waste water on pine bark, Crete 2012 – 3rd international conference on hazardous and industrial waste management, Chania, Crete, September 12-14
- Clark B, Boopathy R (2007), Evaluation of bioremediation methods for the treatment of soil contaminated with explosives in Louisiana Army Ammunition Plant, Minden, Louisiana. *J. Haz. Mat.* 143: 643–648.

- Clausen J.L., Norte N., Dodson M., Robb J. and Rieven S. (2006), *Conceptual Model for the Transport of Energetic Residues from Surface Soil to Groundwater by Range Activities. Final report*. ERDC/CRREL TR-06-18, 155p.
- Ek H., Nilsson E., Birgersson G. and Dave G. (2007), TNT leakage through sediment to water and toxicity to *Nitocra spinipes*. *Ecotoxicology and Environmental Safety*, **67**(3), 341-8.
- EPA 712-C-96-154. (1996), *Ecological Effects Test. Guidelines OPPTS 850.4200 Seed Germination/Root Elongation Toxicity Test*. 6p.
- Esteve-Nunez A., Caballero A. and Ramos J.L. (2001). Biological Degradation of 2,4,6-Trinitrotoluene. *Microbiology and Molecular Biology Reviews*, **65**(3), 335-352.
- Fahrenfeld, N., Zoeckler, J., Widdowson, M.A., Pruden, A. (2012). Effect of biostimulants on 2,4,6-trinitrotoluene (TNT) degradation and bacterial community composition in contaminated aquifer sediment enrichments. *Biodegradation*. In Press.
- Gerth A., Hebner A. and Thomas H. (2003), Natural Remediation of TNT-Contaminated Water and Soil. *Acta Biotechnologica*, **23**(2-3), 143 – 150.
- Green V.S., Stott D.E., Diack M. (2006). Assay for fluorescein diacetate hydrolytic activity: optimization for soil samples. *Soil Biol. Biochem.*, **38**, 693-701.
- Grube M., Muter O., Strikauska S., Gavare M. and Limane B. (2008), Application of FT-IR for control of the medium composition during biodegradation of nitro aromatic compounds. *Journal of Industrial Microbiology and Biotechnology*, **35**, 1545-1549.
- Harrison, I., Vane, C.H. (2010), Attenuation of TNT in seawater microcosms. *Water Science and Technology* **61**(10), 2531-2538.
- Hoffsommer, J. C., Kaplan L. A., Glover D. J., Kubose D. A., Dickinson C., Goya H., Kayser E. G., Groves C. L. and Sitzmann M. E. (1978). *Biodegradability of TNT: A Three-Year Pilot Plant Study*. NSWC/WOL TR 77-136, Naval Surface Weapons Center. Silver Spring, MD.
- Kandeler E. and Gerber H. (1988). Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biology and Fertility of Soils*, **6**, 68–72.
- Kwon, S.-H., (2000), A biological approach in the treatment of TNT wastewater, *Toxicological and Environmental Chemistry*, **75**(1-2), 17-23.
- Lamichhane KM, Babcock Jr RW, Turnbull SJ, Schenck S (2012) Molasses enhanced phyto and bioremediation treatability study of explosives contaminated Hawaiian soils *J. Haz. Mat.* **243**: 334-339.
- Limane B., Juhanson J., Truu J., Truu M., Muter O., Dubova L. and Zarina D. (2009). *Changes in microbial population affected by physico-chemical conditions of soils contaminated by explosives*. In: "Current Research Topics in Applied Microbiology and Microbial Biotechnology", Ed. Antonio Mendez-Vilas World Scientific Publishing Co., 637-640
- Lynch J. C., Brannon J. M. and Delfino J. J. (2002), Effects of Component Interactions on the Aqueous Solubilities and Dissolution Rates of the Explosive Formulations Octal, Composition B, and LX-14. *Journal of Chemical Engineering Data*, **47**, 542–549.
- Lynch, J. C., Myers K. F., Brannon J.M. and Delfino J.J. (2001). Effects of pH and Temperature on the Aqueous Solubility and Dissolution Rate of 2, 4, 6- Trinitrotoluene (TNT), Hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX), and Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (HMX). *Journal of Chemical Engineering Data*, **46**, 1549–1555.
- Martin J. L., Comfort S. D., Shea P. J., Kokjohn T. A. and Drijber R. A. (1997), Denitration of 2,4,6-trinitrotoluene by *Pseudomonas savastanoi*. *Canadian Journal of Microbiology*, **43**, 447–455.
- McCormick, N. G., Feeherry F. E. and Levinson H. S. (1976), Microbial transformation of 2,4,6-TNT and other nitroaromatic compounds. *Applied and Environmental Microbiology*, **31**, 949–958.

- McCutcheon S.C. and Schnoor J.L. (2003), *Phytoremediation: Transformation and Control of Contaminants*, Wiley-IEEE.
- Medina VF, Larson SL, Agwaramgbo L, Perez W, Escalon L (2004) Treatment of trinitrotoluene by crude plant extracts, *Chemosphere* 55: 725–732.
- Meng, Q., Zhao, Q., Zhao, X., Wu, T., Ye, Z. (2012). Removal of Nitro Aromatic Compounds from 2,4,6-Trinitrotoluene Red Water Using 1,2-Ethanediamine Modified Macroporous Polystyrene Microspheres. *Clean - Soil, Air, Water* 40(8), 823-829.
- Mulla SI, Talwar MP, Bagewadi ZK, Hoskeri RS, Ninnekar HZ (2012) Enhanced degradation of 2-nitrotoluene by immobilized cells of *Micrococcus* sp. strain SMN-1, *Chemosphere*. <http://dx.doi.org/10.1016/j.chemosphere.2012.10.030>
- Muter O, Potapova K, Limane B, Sproge K, Jakobsone I, Cepurnieks G, Bartkevics V (2012) The role of nutrients in the biodegradation of 2,4,6-trinitrotoluene in liquid and soil. *J. Environ. Manage.* 98: 51–55.
- Muter O., Versilovskis A., Scherbaka R., Grube M. and Zarina Dz. (2008), Effect of plant extract on the degradation of nitroaromatic compounds by soil microorganisms. *Journal of Industrial Microbiology and Biotechnology*, **35**, 1539-1543
- Nehrenheim, E., Odlare, M. (2010) Treatment of TNT contaminated sludge by using a pilot scale bioreactor – a low cost method for on-site waste management, Second International Conference of Hazardous and Industrial Waste Management, Chania, Crete, Greece, October 5-8, 2010
- Nehrenheim, E., Odlare, M., Allard, B. (2012). Retention of 2,4,6-trinitrotoluene and heavy metals from industrial waste water by using the low cost adsorbent pine bark in a batch experiment. *Water Science and Technology* 64(10), 2052-2058.
- Omstedt A., Edman E., Anderson L.G., Laudon H. (2010), Factors influencing the acid–base (pH) balance in the Baltic Sea: a sensitivity analysis. *Tellus* **62B**, 280–295.
- Pereyra P.J., Rossini G.B., Darrigran G. Toxicity of Neem's oil, a Potential Biocide against the Invasive Mussel *Limnoperna fortunei* (Dunker 1857) *Annals of the Brazilian Academy of Sciences An Acad Bras Cienc* (2012) 84 (4) 1065-1071.
- Phelan J. and Barnett J. L. (2001), Solubility of 2,4-Dinitrotoluene and 2,4,6-Trinitrotoluene in Water. *Journal of Chemical Engineering Data*, **46**, 375–376.
- Qiao H., Wang H., Feng H., Yao J., Shen D. (2010) Reduction and conversion of 2,4,6-trinitrotoluene (TNT) by sulfide under simulated anaerobic conditions, *Journal of Hazardous Materials* 179 (1-3) 989-998
- Ribé, V., Odlare, M., Nehrenheim, E., Riddersand, H., Berglind, R., (2010) Hazard screening by chemical analysis and ecotoxicity bioassays of sediment, ground and surface water sampled from a fire pond and the surrounding area at an explosives manufacturing and processing site, Second International Conference of Hazardous and Industrial Waste Management, Chania, Crete, Greece, October 5-8, 2010
- Ribeiro, E.N., Da Silva, F.T., De Paiva, T.C.B. (2012). Ecotoxicological evaluation of wastewater from 2,4,6-TNT production, *Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering* 47(2), 184-191.
- Rodgers JD, Bunce NJ (2001) Treatment methods for the remediation of nitroaromatic explosives. *Wat. Res.* 35: 2101–2111.
- Rugabber, T.P., Talley, J.W. (2006), Enhancing bioremediation with enzymatic processes: A review. *Practice Periodical of Hazardous, Toxic, and Radioactive Waste Management*, 10(2), 73-85.
- Schackmann A. and Müller R. (1991), Reduction of nitroaromatic compounds by different *Pseudomonas* species under aerobic conditions. *Applied Microbiology and Biotechnology*, **34**, 809–813.

- Schnurer J. and Rosswall T. (1982), Fluorescein Diacetate Hydrolysis as a Measure of Total Microbial Activity in Soil and Litter, *Applied and Environmental Microbiology*, **43**(6), 1256-1261.
- Shen J, He R, Yu H, Wang L, Zhang J, Sun X, Li J, Han W, Xu L (2009), Biodegradation of 2,4,6-trinitrophenol (picric acid) in a biological aerated filter (BAF). *Biores. Technol.* 100: 1922–1930.
- Sikora FJ; Almond RA; Behrends LL; Hoagland JJ; Kelly DA; Phillips WD; Rogers WJ; Summers RK; Thornton FC; Trimm JR; Dader DF (1998), Demonstration results of phytoremediation of explosives-contaminated groundwater using constructed wetlands at the Milan Army Ammunition Plant, Milan, Tennessee Volume IV. Ft. Belvoir Defense Technical Information Center DEC; 394 p.
- Singh, B., Kaur, J., Singh, K. (2012), Microbial remediation of explosive waste (Review). *Critical Reviews in Microbiology* 38(2), 152-167.
- Speitel G. E., Yamamoto H., Autenrieth R. L. and McDonald T. (2002), *Laboratory Fate and Transport Studies of High Explosives at the Massachusetts Military Reservation*. Final Report. University of Texas at Austin and Texas A&M University,
- US Army Environmental Center, (2000), *Pollution Prevention & Environmental Technology Division*, FY-Annual Report, SFIM-AEC-ET-TR-200116
- Wang L.K., Hung Y.-T., Lo H.H. and Yapijakis C (2004), Handbook of industrial and hazardous wastes treatment. 2nd ed. Marcel Dekker Inc.
- Wang Z, Ye Z, Zhang M, Bai X (2010), Degradation of 2,4,6-trinitrotoluene (TNT) by immobilized microorganism-biological filter. *Proc. Biochem.* 45: 993–1001.
- Yasin, M., Shah, A.A., Hameed, A., Ahmed, S., Hasan, F. (2008), Use of microorganisms for the treatment of trinitrotoluene (TNT) containing effluents. *Journal of the Chemical Society of Pakistan*, 30(3), 442-448.
- Zhang, M., Zhao, Q., Ye, Z. (2011), Organic pollutants removal from 2,4,6-trinitrotoluene (TNT) red water using low cost activated coke. *Journal of Environmental Sciences* 23(12), 1962-1969.
- Zhao, Q., Ye, Z., Zhang, M. (2010), Treatment of 2,4,6-trinitrotoluene (TNT) red water by vacuum distillation. *Chemosphere* 80(8), 947-950.